

MD Initials	Emergency Center Orderset for Patients Awaiting Disposition To be initiated when EC patients have been medically cleared and are awaiting disposition	Time	RN Initials
	Secure patient's belongings in labeled "Patient belongings bag"; Security officer to watch until disposition, upon disposition these are to accompany patient to designated facility		
	Diet:		
	Allergies:		
	Vital Signs Every ____ hrs with Neuro checks (may use neuro. flow sheet to document) T: ____ P: ____ BP: ____ RR ____ SaO2 ____ T: ____ P: ____ BP: ____ RR ____ SaO2 ____	_____	
	For extended stay offer pt. hygiene articles/activities of daily living (oral care, shower)		
	☐ Cardiac Monitor ☐ Pulsoximeter (Every ____ hours)		
	PRN Medications:		
	Tylenol (15mg/kg) PO every 4hrs HA\Fever>100.4 F\Pain ____mg given For Fever notify Physician Dr. _____ notified at _____.		
	Motrin (10mg/kg) PO every 6 hrs HA\Fever>100.4 F\Pain ____mg given		
	Maalox\Mylanta 30cc PO every 6 hrs Heartburn\GI upset		
	Haldol ____mg IV\IM\PO every ____ hrs for agitation		
	Ativan ____mg IV\IM\PO every ____ hrs for anxiety/sedation/agitation		
	Benadryl ____mg IV\IM\PO every ____ hrs with Haldol		
	Compazine 10 mg IV\IM\PO every ____ hrs for nausea		
	Other Medications: _____		
	Diabetic Management:		
	Accuchecks: ☐ AC ☐ HS ☐ Every ____ hrs ☐ Other _____ (May document on diabetic flow sheet PRN)		
	Bed Side Blood Glucose _____		
	Lispro (Humalog) insulin subcutaneously using the following sliding scale: BBG less than 70 ----- hypoglycemic procedure BBG 70-120----- give 0 units subcutaneously BBG 121-150 -----give 2 units subcutaneously BBG 151-180 -----give 4 units subcutaneously BBG 181-250----- give 8 units subcutaneously <u>and recheck in 2 hours</u> BBG 251-350-----give 10 units subcutaneously <u>and recheck in 2 hours</u> BBG greater than 350---give 12 units subcutaneously <u>and recheck in 2 hours, notify physician</u> If patient is eating and BBG is greater than 180 on two consecutive BBG checks , increase insulin dose by 2 units at each level. Continue scale at increased level, if still greater than 180 after 8 hours, notify MD regarding uncontrolled hyperglycemia. Consider admission.		
	Social Service consult		
	Neurology consult: Dr. _____ called at (time) _____		
	Psychiatric consult: Dr. _____ called at (time) _____		

RN Signature	RN Initials	RN Signature	RN Initials	MD Signature	MD Initials

T:\Pathways\Patients awaiting Dispo. 4/05, revised 5/05,10/05