

**Table 2. Incubation Periods for Travel-Related Pathogens**

PRIMARY MANIFESTATION	INCUBATION PERIOD		
	SHORT (< 10 DAYS)	INTERMEDIATE (10-21 DAYS)	LONG (> 21 DAYS)
<b>Diarrhea</b>	Campylobacter spp. Enterotoxigenic <i>E. coli</i> Salmonella spp. Shigella spp. Yersinia spp. Rotavirus Noroviruses <i>Vibrio cholera</i>	<i>Cryptosporidium parvum</i> <i>Cyclospora cayetanensis</i> <i>Isospora belli</i> Microsporidian spp.	<i>Dientamoeba fragilis</i> <i>Entamoeba histolytica</i> <i>Giardia intestinalis</i>
<b>Fever</b>	Dengue virus Yellow fever virus <i>Rickettsia africae</i> Nairoviruses Phleboviruses <i>Neisseria meningitidis</i> <i>Pasteurella pestis</i>	Arenavirus West Nile virus <i>Borrelia recurrentis</i> Leptospira spp. <i>Rickettsia typhi</i> <i>Plasmodium falciparum</i>	Hepatitis viruses Hantavirus Rabies virus <i>Coxiella burnetii</i> <i>Plasmodium vivax</i>
<b>Respiratory</b>	Influenza viruses <i>Legionella pneumophila</i> <i>Yersinia pestis</i> <i>Streptococcus pneumoniae</i>	<i>Bordetella pertussis</i>	<i>Paracoccidioides brasiliensis</i> Hantavirus <i>Corynebacterium diphtheriae</i> <i>Mycoplasma pneumoniae</i> <i>Mycobacterium tuberculosis</i>