

Table 2				
Pediatric Dosing of Intravenous Peramivir (Do not exceed 600 mg/day in larger children) <sup>6</sup>				
Age	50-80 mL/min	31-49 mL/min	10-30 mL/min	< 10 mL/min or hemodialysis (on dialysis < days, give post-dialysis)
Birth-30 days	6 mg/kg IV qday	1.5 mg/kg IV qday	1 mg/kg IV qday	0.15 mg/kg IV qday
31 days-90 days	8 mg/kg IV qday	2 mg/kg IV qday	1.3 mg/kg IV qday	0.2 mg/kg IV qday
91 days-180 days	10 mg/kg IV qday	2.5 mg/kg IV qday	1.6 mg/kg IV qday	0.25 mg/kg IV qday
181 days-5 years	12 mg/kg IV qday	3.0 mg/kg IV qday	1.9 mg/kg IV qday	0.3 mg/kg IV qday
6 years-17 years	10 mg/kg IV qday	2.5 mg/kg IV qday	1.6 mg/kg IV qday	0.25 mg/kg IV qday
Creatinine Clearance ( <a href="http://www-users.med.cornell.edu/~spon/picu/calc/crclschw.htm">http://www-users.med.cornell.edu/~spon/picu/calc/crclschw.htm</a> )				