

Table 2. Treatment for von Willebrand's Disease

			Mucous Membrane		Mucous Membrane		Hazardous Areas			
TYPE OF BLEEDING			HEMA-RTHROSIS	DECIDUOUS TEETH						
DIATHESIS	HEMATOMA	HEMATURIA	MOUTH		EPISTAXIS	GASTRO-INTESTINAL	CNS	RETRO-PHARYNGEAL	RETRO-PERITONEAL	
Severe von Willebrand's disease	FFP 10 mL/kg CPT 2 bags/ 10 kg Factor VIII concentrate 10 IU/kg Ice, elastic bandage	Steroids 2 mg/kg x 2 d FFP 10 mL/kg q 12h x 3 d Factor VIII concentrate 10 IU/kg q 12h x 3 d	FFP 5-10 mL/kg CPT 2 bags/ 10 kg Factor VIII concentrate 10 IU/kg	FFP 20 mg/kg Factor VIII concentrate 20 IU/kg once Amicar 100 mg/kg q 6h x 6 d <i>or</i> Cyclokapron 25 mg/kg q 6h x 6 d	Should be seen by MD. If tooth is loose, extract, give replacement therapy as for mouth bleeding	Bilateral pressure on nose for 30 min. Pack with epinephrine-soaked Surgical <i>or</i> Avitene Replacement therapy as for mouth bleeding Amicar 100 mg/kg q 6h x 6 d <i>or</i> Cyclokapron 25 mg/kg q 6h x 6 d	FFP 10 mL/kg q 8h x 3 d CPT 2 bags/ 10 kg q 8h x 3 d Factor VIII concentrate 10 IU/kg q 12h x 6 d	FFP 15 mL/kg q 8h x 3 d CPT 3 bags/ 10 kg q 8h x 3 d Factor VIII concentrate 15 IU/kg q 8h x 5-8 d	Same treatment as hemo-phila A	Same treatment as hemo-phila A
Moderate-mild von Willebrand's disease	Desmopressin (DDAVP) x 4 d	DDAVP x 3 d	DDAVP x 3 d	DDAVP x 3 d with Amicar or Cyclokapron		DDAVP x 5 d		DDAVP x 7 d		

Reprinted with permission from: Hilgartner MW. Factor replacement therapy. In: Hilgartner MW, Pochedly C, eds. *Hemophilia in the Child and Adult*. 3rd ed. New York: Raven; 1989:12-13.